

Excavations, Surveys and Heritage Management in Victoria

Volume 9

2020



Excavations, Surveys and Heritage Management in Victoria
Volume 9, 2020

Edited by

Elizabeth Foley
David Frankel
Susan Lawrence
Caroline Spry

with the assistance of

Ilya Berelov
Shaun Canning

Front cover:

Dead standing black box Culturally Modified Tree along Kromelak (Outlet Creek) (Photo: Darren Griffin)

Excavations, Surveys and Heritage Management in Victoria Volume 9, 2020

Melbourne

© 2020 The authors. All rights reserved.

ISSN 2208-827X

Contents

Editorial note	5
Papers	
Murrup Tamboore: community-led archaeological investigations at the former Keilor Archaeological Area	7
Rebekah Kurpiel, Catherine La Puma, Alex Parmington, Paul Penzo-Kajewski, Ron Jones, Allan Wandin, Bobby Mullins, Nathan Jankowski, Zenobia Jacobs, Molly Thomas, Fleur King and Matthew Meredith-Williams	
‘Scarred for life too’: measuring girth and estimating ages of culturally modified trees using comparative examples from Wotjobaluk Country, Western Victoria	15
Darren Griffin and Abby Cooper	
Results of recent archaeological investigations at Pejark Marsh in Western Victoria	27
Asher Ford, Jocelyn DeJong Strickland and Linda Sonogo	
Investigating anthropogenic and natural disturbance in Aboriginal archaeological contexts: a case study of sand sheets in southeast Melbourne	35
Karen Kapteinis and Caroline Spry	
Update on the Radiocarbon Dating Visualisation Project for Aboriginal places in the State of Victoria, southeastern Australia: insights and issues	51
David Thomas, Caroline Spry, Jacqueline Tumney and Rebekah Kurpiel	
Investigating marsupial bone weathering: implications for understanding the taphonomy of Australian zooarchaeological assemblages	57
Kimberley Crabtree and Jillian Garvey	
Applications of XRD analysis in Australian archaeological contexts: introducing the Olympus TERRA portable XRD analyser	67
Alice Mora	
Cultural heritage significance – not to be muted or trifled with	73
David Tutchener, Rebekah Kurpiel, Bradley Ward, Elizabeth Toohey, Dan Turnbull and Robert Ogden	
A second look at the Langlands Iron Foundry: the engine behind Marvellous Melbourne’s phenomenal rise	79
Sarah Myers, Sarah Mirams and Natalie Paynter	
Going over old ground: modeling historical landscape change in Victoria using GIS	91
Greg Hil, Susan Lawrence and Diana Smith	

Abstracts

An update on research at Berribee Quarry: a dated silcrete extraction site in the Central Murray River Valley of northwestern Victoria	99
Jillian Garvey, Rebekah Kurpiel, Nathan Jankowski, Zenobia Jacobs, Paul Penzo-Kajewski, Darren Perry, Uncle Tinawin Wilson, Bengi Salvi, Austen Graham and Emmy Frost	
Therry (Munro Site) and Elizabeth Street excavations: Site formation along the former Williams Creek, Melbourne CBD	100
Christopher Clark, Sarah Janson, Rebekah Kurpiel, Caroline Spry, Paul Penzo- Kajewski, Brandon Kerrigan, Caroline Hawker, Birgitta Stephenson, Patrick Moss and Sam Player	
Dating the Sunklands; a complex picture	101
Martin Lawler, Aaron Dalla-Vecchia and Jim Wheeler	
Collaboration at the Creek: the Barongarook Creek Bridge Burial, Colac, Victoria	102
Michael Green, Craig Edwards, Owen Cavanough and Matt Grigg	
Indigenous data sovereignty	103
Fiona McConachie, and Renee McAlister	
Hoddle Grid Heritage Review: pre-contact archaeology	104
Petra Schell	
Hoddle Grid Heritage Review: contemporary Aboriginal connections	105
Chris Johnston	
2020 vision: historic artefact management in a new decade	106
Bronwyn Woff and Christine Williamson	
Trashed or treasure: a domestic assemblage from Fitzroy c1860-1880	107
Jennifer Porter and Sharon Lane	
The trouble with MNI – comparing assemblages across Chinese diaspora sites in Australasia	108
Paul Macgregor and Melissa Tsafkas	
A crowd-funded aviation archaeology survey	109
Meaghan L. Aitchison, Leah Byrne, Talia Green, James Kightly and Daniel J. Leahy	
Building a resilient education framework in Victoria archaeology	110
Georgia L. Roberts	
Ten years of OzArch	111
Gary Vines	
10 years of Aboriginal cultural heritage management training: outcomes and prospects	112
Maddy Maitri and Christina Pavlides	

Editorial note

The papers included in this ninth issue of *Excavations, Surveys and Heritage Management in Victoria* were presented at the annual Victorian Archaeology Colloquium held at La Trobe University on 1 February 2020. Once again we had over 150 participants whose attendance testifies to the importance of this fixture within the local archaeological calendar. It continues to be an important opportunity for consultants, academics, managers and Aboriginal community groups to share their common interests in the archaeology and heritage of the State of Victoria.

The papers published here deal with a variety of topics that span Victoria's Aboriginal and European past. While some papers report on the results of specific research projects others focus on aspects of method, approach, education and the social context of our work, and approach.

In addition to the more developed papers, we have continued our practice of publishing the abstracts of other papers given at the Colloquium, illustrated by a selection of the slides taken from the PowerPoint presentations prepared by participants. These demonstrate the range of work being carried out in Victoria, and we hope that many of these will also form the basis of more complete studies in the future. All papers were refereed by the editorial team. This year Elizabeth Foley managed this process and the sub-editing of this volume under the guidance of Caroline Spry. Layout was again undertaken

by David Frankel.

Previous volumes of *Excavations, Surveys and Heritage Management in Victoria* are freely available through La Trobe University's institutional repository, Research Online < www.arrow.latrobe.edu.au:8080/vital/access/manager/Repository/latrobe:41999 >. We hope that this will encourage the dissemination of ideas and information in the broader community, both in Australia and internationally.

We are grateful to the Colloquium's major sponsors ACHM, Ochre Imprints, Ecology and Heritage Partners and Heritage Insight; sponsors Biosis, ArchLink, Christine Williamson Heritage Consultants and Extent; and to La Trobe University for continuing support. We would like to thank them, and all others involved for their generous contributions towards hosting both the event and this publication. Yafit Dahary of 12 Ovens was, as always, responsible for the catering.

Preparation of this volume was, like so much else in 2020, undertaken during the severe restrictions imposed because of the COVID-19 pandemic. We hope that 2021 will be a better year for all and that even if we are unable to hold our Colloquium at the usual time we will be able to do so later in the year.

The editors and authors acknowledge the Traditional Owners of the lands and heritage discussed at the Colloquium and in this volume, and pay their respects to their Elders, past and present.

Applications of XRD analysis in Australian archaeological contexts: introducing the Olympus TERRA portable XRD analyser

Alice Mora

Abstract

X-ray diffraction (XRD) analysis is routinely applied to identify the crystalline phases of a wide range of geological, archaeological, and faunal materials. In recent years, industries have focused on the development of XRD instruments that are increasingly transportable, cost- and time-effective. The quality of data output of portable XRDs is becoming comparable to that of the conventional benchtop XRD systems.

The Olympus TERRA portable XRD analyser features a small vibrating sample holder that requires negligible sample amounts (10–15 mg) in powder form. It is lightweight, battery-operated and can be connected to personal devices via wireless connectivity. Reliable results can be achieved in a short timeframe (5–15 mins). Materials can be quickly analysed on-site with minimal sample destruction. These characteristics make this portable XRD a powerful tool for characterising, identifying and sourcing materials from Australian archaeological contexts.

Introduction

X-ray diffraction (XRD) analysis has proven to be a powerful tool in archaeological studies. This technique is successfully applied to characterise unknown materials and helps archaeologists in reconstructing past human behaviours. Unfortunately, the traditional benchtop XRD systems require a 'significant' amount of material in powder form, extended scanning times and a good level of expertise. These make the application of XRD analysis to archaeological materials challenging, especially in the Australian context, where quarantine regulations and ethical codes of conduct are (for good reasons) in place.

In recent years, industries have developed portable XRD instruments that require minimal sample size, are comfortably transportable, and able to generate inexpensive, quick and reliable data. Among the several portable XRDs available on the market, the

Department of Archaeology and History at La Trobe University has recently acquired the Olympus TERRA portable XRD analyser, whose favourable characteristics will be highlighted in this short paper. The potential applications of XRD analysis in routine archaeological investigations across the Australian landscape will be discussed thereafter, and they include studies aiming to reconstruct past procurement and processing of raw materials, identifying if and how a material may have been heated, or assessing diagenetic alteration of faunal remains.

X-ray diffraction analysis

X-ray powder diffraction analysis is routinely applied to a wide spectrum of geological, archaeological, and faunal materials to identify their constituent crystalline phases (minerals), the presence of amorphous components (e.g. glasses), and the determination of their respective amounts. The rationale behind this technique is that a crystal presents a structure characterised by a distinctive three-dimensional periodic array of atoms, which can diffract X-rays. When the X-rays are focused onto the crystalline phases, they are scattered by the constituting atoms at specific diffraction angles, depending on the periodic nature of a crystalline structure, i.e. the distance between its constituting crystallographic planes (d-spacing). The scattered radiation is collected by a detector, processed, and displayed in a diffractogram. The position, intensity and shape of the diffraction peaks act as a fingerprint for identifying a specific crystal structure (Pecharsky and Zavalij 2005). Therefore, XRD analysis can distinguish between materials that are chemically identical because of the distinctive ordered arrangements of their atoms. A common example is the three distinct forms of silica (SiO₂): glass (amorphous), quartz and cristobalite (both crystalline); which are distinguishable through their diffraction patterns (Smith 1998).

The Olympus TERRA portable XRD analyser

Traditional X-ray diffraction systems have a complex and extensive setup, and their use requires a high level of expertise. In recent years, industries have focused on the

Department of Archaeology and History,
La Trobe University, Bundoora, Vic. 3086
A.Mora@latrobe.edu.au

development of XRD instruments that are increasingly transportable, can be operated after quick training, leading to less costly and time-consuming analyses (Nakai and Abe 2012). The quality of data output of these new portable XRDs is competitive with, if not comparable to that of the conventional benchtop XRD systems.

Among the several companies that have developed portable versions of XRDs, Olympus has placed into the market a very successful product: the 'Olympus TERRA portable XRD analyser' (**Figure 1**). This instrument is very light, weighing around 15 kg with four batteries. It is safely contained in a sturdy box and can be transported on-site within its trolley, making it ideal for use during geological and archaeological fieldwork, as well as in museum and quarantine-regulated collections. This instrument is battery-operated (each set of batteries lasts around 4 hours) and it has wireless connectivity to personal devices (either laptop, smartphone or tablet). The measurement is visualised in real time on the personal device and the output data can be subsequently downloaded. The sample preparation is quick and easy; samples are reduced in particles smaller than 150 μm (100 mesh screen) using the crushing and sieving tools included in the set. The Olympus TERRA pXRD analyser requires only 10-15 mg of sample to run reliable measurements, making the analysis minimally destructive (and feasible in contexts that it would not

normally be). Either bulk or selective sampling can be undertaken, including longitudinal and micro-area sampling.

Conventional XRD setups present a mobile configuration in which the components rotate relative to each other thanks to a goniometer, following Bragg-Brentano geometry. The sample (~ 300 mg) is grounded into a fine powder (<10 μm), homogenised, and appropriately pressed into the sample holder in order to avoid orientation effects. In contrast, in the Olympus TERRA portable XRD, the powder sample is inserted into one of the two sample chambers without any specific preparation thanks to the 'shaker'. This vibrating sample holder ensures that the crystals are randomly oriented by endless grain circulation. The new feature is possible because of the novel transmission geometry, which has first been developed in response to the challenging working conditions of the Mars Curiosity Rover and the limited dimensions of its XRD analyser (CheMin) (Downs and MSL Science Team 2015). In this type of configuration, the components are in a fixed position: the X-rays leave the tube (Cu or Co target), pass through a collimator, and collide into the sample, where they are diffracted by the array of grains onto a Charge-Coupled Device (CCD) detector. This allows the creation of an instrument of limited dimensions, easily transportable, which requires only minimal maintenance. Since the CCD camera can collect both diffraction and fluorescence data, the Olympus TERRA pXRD is at the same time an X-ray diffractometer and X-ray spectrometer (with an XRF energy range of 2.5 to 25 keV).

Another significant advantage is that the complete diffraction pattern of the sample is collected and displayed simultaneously (over the entire angular range of 5° to 55°) after the first exposure (~ 20 sec). In addition, the fluorescence data from the same sample is concurrently visualised in an XRF spectrum. This is adequate to quickly screen samples, such as understanding whether a known crystalline phase is present in the material, or to discriminate between two possible phases. Reliable data output can be achieved in less than 15 minutes, after approximately 50 exposures. More exposures will sharpen the diffraction peaks and reduce the background noise (**Figure 2**). The XRD data produced by the Olympus TERRA portable XRD analyser can be treated using two software packages: X Powder Software for Qualitative and Semi-Quantitative analysis, and Siroquant Software for Quantitative Rietveld analysis.

When using a conventional benchtop XRD, it takes around 10 minutes to quickly acquire the complete diffraction pattern because the sample is scanned at every possible angle. The resulting diffractogram will present a wider angular range ($>55^\circ$) and a slightly better detection limit (1-2 wt%) than that produced by a pXRD such as the Olympus TERRA. As outlined earlier, both the



Figure 1. The Olympus TERRA portable XRD analyser and the 'shaker'




Figure 2. XRD diffractograms of emu (*Dromaius novaehollandiae*) eggshell (calcite) and freshwater bivalve (*Alathyria jacksoni*) shell (aragonite, calcite, quartz), using an Olympus TERRA pXRD analyser

benchtop and portable XRD systems present distinctive benefits and drawbacks, and ultimately the choice of using one of the two depends on several variables, including the research questions, available funds and time, level of expertise, and sampling constraints (e.g. sample preservation, sample location, regulations). As a rule of thumb, the portable XRD analyser would be efficient for a quick (on-site) screening of all the samples (with identification and quantification of all minerals), and for selecting the more complex mixtures that could be subsequently analysed via a benchtop XRD system.

XRD applications in Australian archaeological contexts

Broadly speaking, X-ray diffraction analysis identifies the mineralogical composition of materials, permitting an understanding of the conditions under which each material was formed and subsequently altered. XRD analysis presents a myriad of potential applications in Australian archaeological contexts, depending on specific research questions and type of samples, which

may include faunal remains, lithic materials, ceramics, heat-retainers, ochres, and sediments.

Archaeofaunal remains such as fragments of bones, teeth, mollusc shell and eggshell are commonly found across the Australian landscape. The inorganic fraction of both bones and teeth is constituted by hydroxylapatite crystals. By studying the microstructural alterations of bone/tooth mineral crystallites, it may be possible to identify diagenetic and thermal processes that affected the faunal material (Piga et al. 2009; Rahmat et al. 2020; Rogers et al. 2010), which would help reconstructing ancient fire use and food processing (Solari et al. 2015; Van Hoesel et al. 2019), and/or post-mortem environmental conditions to which they were exposed (Stathopoulou et al. 2008; Trueman et al. 2004; Tütken et al. 2008).

Calcite, the inorganic fraction of avian eggshell, is also affected by heat-induced mineralogical changes, which are dependent on the temperature of heating (Engin et al. 2006; Macha et al. 2015; Naemchanthara et al. 2008; Tsuboi and Koga 2018). XRD analysis of mollusc shells and fish otoliths allows the identification of the different forms of calcium carbonate (aragonite and/or calcite) constituting their mineral fractions, which are indicative of the environmental conditions during the life of the animal and post-mortem. Environmental factors, such as chemical composition of water during shell formation, may affect the resulting shell mineralogy (Checa et al. 2007; Medaković et al. 2003). Moreover, mineralogical transition of aragonite into calcite may be later induced by heating as a result of food processing (Aldeias et al. 2019) or post-depositional thermal alteration (Milano and Nehrke 2018). The mineralogical characterization of the archaeofaunal remains is also a powerful tool for selecting well-preserved and reliable samples for DNA studies (Götherström et al. 2002), stable isotope analysis (Disspain et al. 2016; Munro et al. 2007), and radiocarbon dating (Long et al. 2018; Webb et al. 2007).

The determination of the mineralogical composition of lithic materials and earth-based pigments may help establish the provenance of raw materials by identifying the possible geological sources, and thus giving insight into procurement strategies and histories of transport and use (Corkill 2005; Dayet et al. 2016; Jercher et al. 1998; Trindade et al. 2010). Characterising the mineralogy of ochres may also solve attribution and authenticity issues of Indigenous artworks (Nel et al. 2010). The assessment of mineralogical changes in ceramics, heat-retainers, and other burnt clays allows for the reconstruction of their thermal history including the determination of firing events and maximum temperatures (Holakooei et al. 2014; Rasmussen et al. 2012). Clayey sediments and soils exposed to fire (hearths) undergo mineralogical alterations, which can be indicative of human occupation of the site and fire-related activities (Berna et al. 2007; Singh et al. 1991).

To conclude, the Olympus TERRA pXRD analyser may be a cost- and time-effective tool for characterizing, identifying, and sourcing (on-site and with minimal sample destruction) a range of archaeological materials that are routinely found in Australian contexts. Given its practicality, the use of a portable XRD could easily be implemented in routine archaeological investigations. If needed, more complex studies could subsequently be undertaken via a benchtop XRD system.

Acknowledgments

I would like to thank the anonymous reviewers for their helpful suggestions.

References

- Aldeias, V., S. Gur-Arieh, R. Maria, P. Monteiro and P. Cura 2019 Shell we cook it? An experimental approach to the microarchaeological record of shellfish roasting. *Archaeological and Anthropological Sciences* 11(2):389–407
- Berna, F., A. Behar, R. Shahack-Gross, J. Berg, E. Boaretto, A. Gilboa, I. Sharon, S. Shalev, S. Shilstein, N. Yahalom-Mack, J.R. Zorn, and S. Weiner 2007 Sediments exposed to high temperatures: Reconstructing pyrotechnological processes in Late Bronze and Iron Age Strata at Tel Dor (Israel). *Journal of Archaeological Science* 34(3):358–373
- Checa, A.G., C. Jiménez-López, A. Rodríguez-Navarro and J.P. Machado 2007 Precipitation of aragonite by calcitic bivalves in Mg-enriched marine waters. *Marine Biology* 150(5):819–827
- Corkill, T. 2005 Sourcing stone from the Sydney region: A hatchet job. *Australian Archaeology* 60(1):41–50
- Dayet, L., F.-X. Le Bourdonnec, F. Daniel, G. Porraz and P.-J. Texier 2016 Ochre provenance and procurement strategies during the middle stone age at Diepkloof Rock Shelter, South Africa. *Archaeometry* 58(5):807–829
- Disspain, M.C.F., S. Ulm, C. Izzo and B.M. Gillanders 2016 Do fish remains provide reliable palaeoenvironmental records? An examination of the effects of cooking on the morphology and chemistry of fish otoliths, vertebrae and scales. *Journal of Archaeological Science* 74:45–59
- Downs, R.T. and MSL Science Team 2015 Determining mineralogy on Mars with the CheMin X-ray diffractometer. *Elements* 11(1):45–50
- Engin, B., H. Demirtaş and M. Eken 2006 Temperature effects on egg shells investigated by XRD, IR and ESR techniques. *Radiation Physics and Chemistry* 75(2):268–277
- Götherström, A., M.J. Collins, A. Angerbjörn and K. Lidén 2002 Bone preservation and DNA amplification. *Archaeometry* 44(3):395–404
- Holakooei, P., U. Tessari, M. Verde and C. Vaccaro 2014 A new look at XRD patterns of archaeological ceramic bodies. *Journal of Thermal Analysis and Calorimetry* 118(1):165–176
- Jercher, M., A. Pring, P.G. Jones and M.D. Raven 1998 Rietveld X-ray diffraction and X-ray fluorescence analysis of Australian Aboriginal ochres. *Archaeometry* 40(2):383–401
- Long, K., R. Wood, I.S. Williams, J. Kalish, W. Shawcross, N. Stern and R. Grün 2018 Fish otolith microchemistry: Snapshots of lake conditions during early human occupation of Lake Mungo, Australia. *Quaternary International* 463:29–43
- Macha, I.J., L.S. Ozyegin, F.N. Oktar and B. Ben-Nissan 2015 Conversion of ostrich eggshells (*Struthio camelus*) to calcium phosphates. *Journal of The Australian Ceramic Society* 51(1):125–133
- Medaković, D., R. Slapnik, S. Popović and B. Gržeta 2003 Mineralogy of shells from two freshwater snails *Belgrandiella fontinalis* and *B. kuesteri*. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* 134(1):121–127
- Milano, S. and G. Nehrke 2018 Microstructures in relation to temperature-induced aragonite-to-calcite transformation in the marine gastropod *Phorcus turbinatus*. *PloS one* 13(10):e0204577
- Munro, L.E., F.J. Longstaffe and C.D. White 2007 Burning and boiling of modern deer bone: Effects on crystallinity and oxygen isotope composition of bioapatite phosphate. *Palaeogeography, Palaeoclimatology, Palaeoecology* 249(1–2):90–102
- Naemchanthara, K., S. Meejoo, W. Onreabroy and P. Limsuwan 2008 Temperature effect on chicken egg shell investigated by XRD, TGA and FTIR. *Advanced Materials Research* 55–57:333–336
- Nakai, I. and Y. Abe 2012 Portable X-ray powder diffractometer for the analysis of art and archaeological materials. *Applied Physics A* 106(2):279–293
- Nel, P., P.A. Lynch, J.S. Laird, H.M. Casey, L.J. Goodall, C.G. Ryan and R.J. Sloggett 2010 Elemental and mineralogical study of earth-based pigments using particle induced X-ray emission and X-ray diffraction. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* 619(1–3):306–310
- Pecharsky, V. and P. Zavalij 2005 Fundamentals of diffraction. In V. Pecharsky and P. Zavalij (eds), *Fundamentals of powder diffraction and structural characterization of materials*, pp. 99–260. USA: Springer Science+Business Media
- Piga, G., T.J.U. Thompson, A. Malgosa and S. Enzo 2009 The potential of X-ray diffraction in the analysis of burned remains from forensic contexts. *Journal of Forensic Sciences* 54(3):534–539
- Rahmat, R.A., M.A. Humphries, J.J. Austin, A.M.T. Linacre, M. Raven and P. Self 2020 Integrating spectrophotometric and XRD analyses in the investigation of burned dental remains. *Forensic Science International* 310:110236

- Rasmussen, K.L., G.A. De La Fuente, A.D. Bond, K.K. Mathiesen and S.D. Vera 2012 Pottery firing temperatures: A new method for determining the firing temperature of ceramics and burnt clay. *Journal of Archaeological Science* 39(6):1705–1716
- Rogers, K., S. Beckett, S. Kuhn, A. Chamberlain and J. Clement 2010 Contrasting the crystallinity indicators of heated and diagenetically altered bone mineral. *Palaeogeography, Palaeoclimatology, Palaeoecology* 296(1–2):125–129
- Singh, B., S. O'Connor, P. Veth and R. Gilkes 1991 Detection of amorphous aluminosilicate by X-ray diffraction and chemical analysis to detect firing in archaeological sediments. *Archaeology in Oceania* 26(1):17–20
- Smith, D.K. 1998 Opal, cristobalite, and tridymite: Noncrystallinity versus crystallinity, nomenclature of the silica minerals and bibliography. *Powder Diffraction* 13(1):2–19
- Solari, A., D. Olivera, I. Gordillo, P. Bosch, G. Fetter, V.H. Lara and O. Novelo 2015 Cooked bones? Method and practice for identifying bones treated at low temperature. *International Journal of Osteoarchaeology* 25(4):426–440
- Stathopoulou, E.T., V. Psycharis, G.D. Chryssikos, V. Gionis and G. Theodorou 2008 Bone diagenesis: New data from infrared spectroscopy and X-ray diffraction. *Palaeogeography, Palaeoclimatology, Palaeoecology* 266(3–4):168–174
- Trindade, M.J., M.I. Dias, J. Coroado and F. Rocha 2010 Firing tests on clay-rich raw materials from the Algarve basin (southern Portugal): Study of mineral transformations with temperature. *Clays and clay minerals* 58(2):188–204
- Trueman, C.N., A.K. Behrensmeyer, N. Tuross and S. Weiner 2004 Mineralogical and compositional changes in bones exposed on soil surfaces in Amboseli National Park, Kenya: Diagenetic mechanisms and the role of sediment pore fluids. *Journal of Archaeological Science* 31(6):721–739
- Tsuboi, Y. and N. Koga 2018 Thermal decomposition of biomineralized calcium carbonate: Correlation between the thermal behavior and structural characteristics of avian eggshell. *ACS Sustainable Chemistry & Engineering* 6(4):5283–5295
- Tütken, T., T.W. Vennemann and H.-U. Pfretzschner 2008 Early diagenesis of bone and tooth apatite in fluvial and marine settings: Constraints from combined oxygen isotope, nitrogen and REE analysis. *Palaeogeography, Palaeoclimatology, Palaeoecology* 266(3–4):254–268
- Van Hoesel, A., F.H. Reidsma, B.J.H. van Os, L. Megens and F. Braadbaart 2019 Combusted bone: Physical and chemical changes of bone during laboratory simulated heating under oxidising conditions and their relevance for the study of ancient fire use. *Journal of Archaeological Science: Reports* 28:102033
- Webb, G.E., G.J. Price, L.D. Nothdurft, L. Deer and L. Rintoul 2007 Cryptic meteoric diagenesis in freshwater bivalves: Implications for radiocarbon dating. *Geology* 35(9):803–806